

Mid-Infrared Images of Stars and Dust in Irregular Galaxies¹

Deidre A. Hunter², Bruce G. Elmegreen³, and Emily Martin^{2,4}

ABSTRACT

We present mid-infrared to optical properties of 22 representative irregular galaxies: 18 Im, 3 BCDs, and one Sm. The mid-IR is based on images from the *Spitzer Space Telescope* archives. The 3.6 and 4.5 μm bands and the *UBVJHK* images are used to examine disk morphology and the integrated and azimuthally averaged magnitudes and colors of stars. The non-stellar contribution to the 4.5 μm images is used to trace hot dust. The 5.8 and 8.0 μm images reveal emission from hot dust and PAHs, and both may contribute to these passbands, although we refer to the non-stellar emission as PAH emission. We compare the 8.0 μm images to $\text{H}\alpha$. Im galaxies have no hidden bars, and those with double-exponential optical light profiles have the same at mid-IR. Most galaxies have similar optical and mid-IR scale lengths. Four galaxies have super star clusters that are not visible at optical bands. Galaxies with higher area-normalized star formation rates have more dust and PAH emission relative to starlight. Hot dust and PAH emission is found mostly in high surface brightness H II regions, implying that massive stars are the primary source of heating. Galaxies with intense, wide-spread star formation have more extended PAH emission. The ratio of PAH to $\text{H}\alpha$ emission is not constant on small scales. PAHs are associated with shells and giant filaments, so they are not destroyed during shell formation.

Subject headings: galaxies: irregular — galaxies: ISM — galaxies: photometry — galaxies: stellar content — galaxies: structure

1. Introduction

The *Spitzer Space Telescope* (*Spitzer*) allows us a deeper view into the infrared emission of galaxies than was possible with previous infrared satellites. The InfraRed Array Camera

¹This work is based in part on archival data obtained with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA.

²Lowell Observatory, 1400 West Mars Hill Road, Flagstaff, Arizona 86001 USA

³IBM T. J. Watson Research Center, PO Box 218, Yorktown Heights, New York 10598 USA

⁴Current address Wheaton College, Norton MA 02766

(IRAC; Fazio et al. 2004) in particular gives us the opportunity to image galaxies in pass-bands that capture the light of stars and emission from hot dust and polycyclic aromatic hydrocarbons (PAHs) with a spatial resolution that is comparable to that of ground-based optical images. IRAC’s Channel 1 ($3.6\ \mu\text{m}$) centered at $3.6\ \mu\text{m}$ probes the stellar population; Channel 2 ($4.5\ \mu\text{m}$) centered at $4.5\ \mu\text{m}$ contains emission from hot dust as well as stars; and Channel 3 ($5.8\ \mu\text{m}$) at $5.8\ \mu\text{m}$ and Channel 4 ($8.0\ \mu\text{m}$) at $7.8\ \mu\text{m}$ are dominated by thermal emission from PAHs in the interstellar medium (ISM) along with emission from hot dust. The $8.0\ \mu\text{m}$ image in particular contains the 7.7 and $8.6\ \mu\text{m}$ emission bands due to PAHs, and the $5.8\ \mu\text{m}$ image contains the $6.2\ \mu\text{m}$ band (Léger & Puget 1984, Allamandola et al. 1989).

Irregular (Im) galaxies, the most numerous type of galaxy in the universe, have received limited attention with infrared satellite telescopes because of their faintness which makes observations difficult. However, they are an interesting class to examine in the infrared because of their distinct differences compared to giant spiral systems. As a class the Im galaxies have a lower dust content, and their molecular clouds are, to a greater extent than spirals, the cores of atomic clouds. They also are generally dominated in the optical by their young stellar populations. They are physically smaller, lower in luminosity and surface brightness, and more gas-rich. Most Im galaxies are forming stars; yet they do so without the benefit of spiral density waves. The Im galaxies may represent the nature of star formation in the early universe.

Generally, PAH emission is low in dwarf galaxies, and this may be related to the low metallicity and intense stellar radiation field in these systems relative to those of spirals (Boselli et al. 1998, Thuan et al. 1999, Madden 2000, Sturm et al. 2000, Galliano et al. 2003, Lu et al. 2003, Houck et al. 2004, Rosenberg et al. 2006). Hunter et al. (2005), for example, used *Infrared Space Observatory* mid-infrared imaging ($6.75\ \mu\text{m}$ and $15\ \mu\text{m}$) and far-infrared (FIR) spectroscopy to examine the properties of five Im systems. They found that the PAH emission, which is associated only with the brightest H II regions, is depressed relative to that of small grains, the FIR, and H α . In addition, the integrated [CII] emission relative to PAH emission is unusually high compared to that in spirals. These two observations suggest that Carbon is more likely to be in an atomic form than a PAH. Recently, in a study of Sloan Digital Sky Survey galaxies observed in the IRAC First Look Survey, Hogg et al. (2005) found that low luminosity blue galaxies have anomalously low PAH emission relative to starlight. Similarly, Engelbracht et al. (2005) have examined a sample of low metallicity systems, and found that the $8.0\ \mu\text{m}$ -band nebular emission decreases abruptly relative to the $24\ \mu\text{m}$ emission from dust for galaxies with metallicities less than $1/3$ – $1/5$ solar. As Draine (2005) has pointed out, the ISM of low metallicity systems seems to be a more hostile environment for PAHs for reasons that are not entirely clear.

Here we use the unique capabilities of the IRAC camera to further our understanding of the stars and dust in dwarf galaxies through an investigation of 22 irregular galaxies in the mid-IR. We examine the distribution and intensity of hot dust and PAH emission in detail and compare them to that of $H\alpha$, a tracer of the current star formation activity. We also probe the stellar populations with the $3.6\ \mu\text{m}$ and $4.5\ \mu\text{m}$ images, determining the stellar distributions and colors and comparing the mid-IR to the optical.

The sample includes 18 Im systems, 3 blue compact dwarfs (BCDs), and one Magellanic-type spiral (Sm) galaxy. Except for NGC 4449, these are systems included in the $H\alpha$ and broad-band survey of Im/BCD/Sm galaxies conducted by Hunter & Elmegreen (2004, 2006). The galaxies and some pertinent parameters are listed in Table 1. The galaxies included in this study span a large range in properties: M_V ranges from -11.7 to -18.3 ; the star formation rate ranges from no detectable $H\alpha$ emission (M81dwA and DDO 210) to galaxies undergoing a burst of star formation (IC 10, NGC 1705). This sample, although defined by what was available in the *Spitzer* archives, is fairly representative of the Im class and is not dominated by high star formation rate dwarf galaxies which have been the subject of most past studies of dwarfs in the infrared. Furthermore, except for NGC 4449, these are relatively isolated systems; the galaxies are not involved in an obvious on-going interaction with another galaxy.

2. Data Reduction and Analysis

The galaxies were imaged with the IRAC detectors; 16 of the galaxies were imaged in all four pass-bands and 6 were imaged only at $4.5\ \mu\text{m}$ and $8.0\ \mu\text{m}$. The imaging employed various dithering patterns, and large galaxies were mapped. All galaxies had short and long or short, medium, and long exposures. Many of the galaxies were observed in two separate observing sessions (those from the SINGS Legacy database, Kennicutt et al. 2003).

The data were obtained from the *Spitzer* archive and reduced with the Basic Calibrated Data pipeline. Post-pipeline reductions were done using the *Spitzer* contributed software package MOPEX. Saturated pixels in the long exposures were replaced with appropriate values determined from the short or medium exposures. Overlapping fields were placed onto a single grid determined to include all frames for each galaxy, and backgrounds were matched. Optical distortion was removed and the images were repixelated to $1.2''$ per pixel. We used dual-outlier and mosaic-outlier detection algorithms to remove cosmic rays, and experimented with different outlier rejection parameters before settling on thresholds and other parameters that best removed the cosmic rays and bad pixels without removing real objects. We did not do corrections for “muxbleed” or “column pull-down” problems since

the corrections are not adequate for photometry; we preferred to be able to identify these flaws in the final images so that they could be masked and excluded in the photometry.

We determined the background in the images themselves from regions not containing galaxy. In some cases the background was sufficiently flat that a simple constant was used. In other cases, we did a two-dimensional fit, usually of order two, to the background. The background was subtracted from the image.

To prepare the images for photometry, we needed to remove the foreground stars and background galaxies contaminating the object galaxy image, as we had done for the UB-VJHK images (Hunter & Elmegreen 2006). We used the V -band image of the galaxy and surrounding non-galaxy regions in the IRAC images as guides to what was likely to be galactic. In most cases we edited objects from the image by interpolating across a circle centered on the star to be removed. The editing was done star by star for the $3.6\ \mu\text{m}$ image, and the commands were logged to a file; then the logged commands were applied to the $4.5\ \mu\text{m}$ image so that the $3.6\ \mu\text{m}$ and $4.5\ \mu\text{m}$ images were cleaned in an identical manner.

In some cases this method of removing foreground stars was not feasible (DDO 69, DDO 75, DDO 216, IC 10, NGC 3109, NGC 4214, and NGC 6822) either because the galaxy itself was highly resolved or the foreground contamination was particularly high. In those cases we used the region surrounding the galaxy to determine a constant background that included a statistical sampling of foreground stars. We tried both techniques on a few galaxies, particularly those in which the slopes in the exponential disks were different in $3.6\ \mu\text{m}$ than in V . We found that the choice of technique for foreground subtraction did not make any significant difference to the photometry.

In NGC 6822, Milky Way cirrus was prevalent in the $8.0\ \mu\text{m}$ image. Thus, we did not extract quantitative information, but we did make a qualitative spatial comparison with the $\text{H}\alpha$ image, restricting ourselves to the higher surface brightness emission.

The IRAC images were flux calibrated and pixel values converted to units of MJy sr^{-1} in the Basic Calibrated Data pipeline. We express quantities here as magnitudes relative to Vega where we have used fluxes of 280.9, 179.7, 115.0, and 64.1 Jy for Vega for bands $3.6\ \mu\text{m}$, $4.5\ \mu\text{m}$, $5.8\ \mu\text{m}$, and $8.0\ \mu\text{m}$, respectively (Reach et al. 2005), and we will refer to these images and quantities measured from them as [3.6], [4.5], [5.8], and [8.0], respectively. The Johnson UBV photometric system is also normalized to Vega, so the IR-optical combination of $V - [3.6]$ that we will use is in terms of the Vega IR-optical color. The calibration is based on point sources with photometry performed through a $12''$ radius aperture. An aperture correction to extended sources is, therefore, necessary. The corrections we adopted are 0.063, 0.071, 0.281, and 0.331 mag for [3.6], [4.5], [5.8], and [8.0], respectively, from Reach et al.

(2005).

No reddening corrections have been applied to the IRAC photometry. Internal reddening in irregular galaxies is low (Hunter et al. 1986, 1989), and in most cases, the foreground reddening is zero or near zero.

The [4.5] passband contains emission from stars and hot dust, and the [5.8] and [8.0] passbands include PAH emission in addition to starlight. We will refer to the hot dust emission in [4.5] as “[4.5]_{dust}” and that from the stars as “[4.5]_{*}.” The [5.8] and [8.0] images also contain emission from warm dust (Rosenberg et al. 2006). PAH emission-bands are found in the spectra of Im galaxies, so emission from both PAHs and dust could, in principle be present (Engelbracht et al. 2005). We will usually refer to the non-stellar emission in [8.0] as “PAH emission” for simplicity in order to distinguish it from stellar emission, but keep in mind that both PAHs and hot dust may contribute to the [5.8] and [8.0] passbands. All quantities measured from [5.8] and [8.0] are measured from the star-subtracted images, and are referred to as “[5.8]_{PAH}” and “[8.0]_{PAH}”.

We have used the [3.6] images to remove the stars from the other passbands and leave non-stellar emission. To do this, we used foreground stars in each image to determine a scaling factor that matches the stellar photometry in the [3.6] image to that in the other images, scaled the [3.6] image, and subtracted the scaled [3.6] image. The scaling was done empirically, so no explicit assumption was made concerning the spectral type of the stars. In the case of [4.5], this sometimes resulted in an oversubtraction of the galaxy (IC 10, NGC 3738, NGC 4449), presumably because the galaxy is blue and the foreground stars are red. In those cases we empirically adjusted the scaling factor by 5–11% to make the stars go away without oversubtracting the galaxy itself. The average factor that the 3.6 μm images were divided by were 1.59 ± 0.08 , 2.44 ± 0.18 , and 4.23 ± 0.30 for the 4.5 μm , 5.8 μm , and 8.0 μm images, respectively. The uncertainty is the variation between galaxy fields. For those galaxies with only 4.5 μm and 8.0 μm images, the 4.5 μm image was divided by 2.72 ± 0.14 , on average, before being used to remove stars from the 8.0 μm image. For those galaxies as well, a [4.5]_{dust} image could not be constructed and the original [4.5] image was taken to be [4.5]_{*}.

Having constructed a hot dust image as just described, we subtracted the hot dust from the original [4.5] image to leave an image of only starlight in this passband. According to Regan et al. (2004) we should expect most of the emission in the [4.5] image to be starlight for galaxies undergoing quiescent star formation, and indeed, the contribution from hot dust was small in our systems.

In the highly resolved galaxy NGC 6822 we found that there were stars left in the [4.5]_{dust}

and $[5.8]_{\text{PAH}}$ images after scaling and subtracting the $[3.6]$ image. These stars clustered around NGC 6822, so they are likely stars in NGC 6822 rather than foreground objects. We were curious as to the nature of these stars. Therefore, we measured brightnesses of those stars as well as of a sample of NGC 6822 stars that did subtract cleanly. We did the photometry on the IRAC images as well as on JHK 2MASS images that we extracted from the 2MASS archive. We compared the location of the stars on the $J-H$ vs $H-K$ color-color diagram with stellar photometry of the LMC presented by Nikolaev & Weinberg (2000). Stars that had subtracted cleanly lie in the region of the color-color diagram populated by dwarf and giant stars. Stars that did not subtract successfully lie where one expects Carbon stars and B[e] stars to sit. Thus, it is likely that the stars that remained in the $[4.5]_{\text{dust}}$ and $[5.8]_{\text{PAH}}$ images after scaling and subtracting the $[3.6]$ image are real objects in NGC 6822. To construct the non-stellar dust and PAH images, these stars were edited out of the $[4.5]_{\text{dust}}$ and $[5.8]_{\text{PAH}}$ images by hand.

For most of our sample galaxies, we have $H\alpha$ and UBV images, and for a few we also have JHK images (Hunter & Elmegreen 2004, 2006). In order to compare the galaxies’ emission in these different passbands, we geometrically transformed all images to match the scale and orientation of the V -band image. Surface photometry was extracted in ellipses using the parameters—center, ellipticity, major axis step size—that were used for photometry of the V -band image. The photometry parameters for all galaxies except NGC 4449 are listed by Hunter & Elmegreen (2006). The original optical surface photometry for NGC 4449 was presented by Hunter et al. (1999) who used a position angle and ellipticity that changed with radius because of the substantial stellar bar. Here we have used the position angle (64°) and ellipticity ($b/a = 0.73$) that are appropriate for the outer disk to be consistent with our other photometry, and have redone the optical surface photometry to match.

In the ellipse surface photometry, bad pixels were masked and replaced with an average of the brightness in the annulus. Some galaxies were not completely covered on one or more sides by the IRAC maps (IC 10, NGC 4214, NGC 4449). The portion of the galaxy that was blank was treated as for masked pixels, and the surface photometry was truncated when the portion that was missing became larger than the portion with measurable galaxy. The IRAC image of NGC 6822 has a field of view that is large enough to include the entire galaxy, whereas our optical image does not. As a result, the IRAC surface photometry only goes as far out in radius as the optical goes, but for integrated IRAC values we used the original IRAC image and its larger field of view.

The uncertainties in the surface photometry followed the formula given by Hunter & Elmegreen (2006). The original sky background was reconstructed for this purpose using the frame duration time and estimates of the zodiacal and Galactic backgrounds (FRAMTIME,

ZODY_EST, and ISM_EST key words in the header of the images).

The galaxies are shown in Figure 1. There are potentially 3 images, all showing the same field of view, and two graphs for each galaxy. These are arranged in two rows per galaxy and two galaxies per page. The upper left image for each galaxy is the geometrically transformed [3.6] image, or logarithm of [3.6] if the galaxy is high in surface brightness. An outer V -band contour is plotted on this image for comparison. For the galaxies with no [3.6] image, the [4.5] image is shown instead. The upper middle image is an $H\alpha$ image with a contour of the [8.0]_{PAH} PAH emission image superposed. The lower left image is the same but contours are of the [4.5]_{dust} image. If the galaxy has no detectable hot dust, this image is left blank. If the galaxy has no detectable PAH emission, the $H\alpha$ image is shown with the same outer V -band contour superposed that was used on the [3.6] or [4.5] image. We do not show $H\alpha$ images for M81dwA and DDO 210 because no $H\alpha$ emission was detected in these galaxies. In those cases there is also no hot dust or [8.0]_{PAH} PAH emission, so nothing is shown in the upper middle and bottom left spots. The graph in the bottom middle spot is the surface photometry in V , J if observed, $H\alpha$, [3.6] or [4.5]_{*}, and [8.0]_{PAH} PAH emission, if detected, as a function of radius. The solid lines are fits to the exponential disks in V , J , and [3.6] or [4.5]_{*}. The graph on the right is a plot of the colors as a function of radius.

Integrated photometry is given in Table 2 along with the central surface brightness $\mu_0^{[3.6]}$ and disk scale length $R_D^{[3.6]}$ determined from the exponential fit to the surface photometry for the galaxies with observations in all 4 IRAC passbands. Similar quantities are given in Table 3 for the galaxies with observations in only [4.5] and [8.0] passbands. Upper limits are $5\times\text{rms}$ over an area of radius 3 pixels; the rms is determined from sampling across the image.

3. Stars

3.1. Integrated properties

The [3.6] image offers an alternate perspective on the stellar populations of a galaxy compared to the optical. In principle the near-IR is dominated by old stars and low mass stars of all ages, although young red supergiants also contribute from pockets of recent star formation. The integrated galactic [3.6] magnitudes are reported in Table 2 as an absolute magnitude $M_{[3.6]}$ using the distances in Table 1. Several integrated colors and M_V are plotted in Figure 2 as a function of $M_{[3.6]}$. There is a strong correlation between the two absolute magnitudes M_V and $M_{[3.6]}$, as one would expect. In terms of the colors there are correlations between $M_{[3.6]}$ and $(B-V)_0$ and between $M_{[3.6]}$ and $V-[3.6]$, and these correlations are in the

sense that more luminous galaxies are redder. The change in $(B-V)_0$ is not large, however; over 8 mag in $M_{[3.6]}$ $(B-V)_0$ reddens by only ~ 0.2 mag and the scatter is comparable to the change. The change in $V-[3.6]$ is larger; $V-[3.6]$ changes by 1.25 mag over that same interval.

The color $[3.6]-[4.5]_*$, on the other hand, is constant among most of the galaxies at an average value of -0.08 ± 0.18 . This color is comparable to that of an early K star (Pahre et al. 2004, Willner et al. 2004, Reach et al. 2005). In a study of a sample of galaxies spanning the Hubble sequence, Pahre et al. (2004) found that $[3.6]-[4.5]$ becomes redder with later Hubble type. The change is small, only ~ 0.15 mag from early to late-type galaxies, but it reaches about ~ 0 at $T = 10$ (Im galaxies). The small difference between their $[3.6]-[4.5]$ color and ours for Im galaxies could be due to the fact that their $[4.5]$ magnitude includes the hot dust component. Therefore, their $[3.6]-[4.5]$ color should be a little bit redder than ours, as is observed. Furthermore, the change in $[3.6]-[4.5]$ color that they observe could be due, at least in part, to the increasing contribution of hot dust in later type galaxies. The average color of our sample is similar to that observed for the disk of M81 by Willner et al. (2004; see their Figure 2).

The fact that the integrated stellar color $[3.6]-[4.5]_*$ is constant from galaxy to galaxy suggests simply that the old and low mass stellar component of galaxies have nearly the same colors. By contrast, that $V-[3.6]$ changes more steeply with galactic absolute magnitude is an indication of the relative importance of the young stellar component to the older, low mass component. Irregular galaxies are blue in optical colors primarily because of the dominance of the young stellar population (and partly due to low metallicity). Therefore, it makes sense that $V-[3.6]$ should reflect this.

3.2. Large-scale Morphology

In some spirals, stellar structures, such as bars, become more recognizable in the infrared, perhaps because they are formed from old stars or dust obscures their features in the optical. In our galaxy sample, however, the large-scale morphology that we see in the $[3.6]$ or $[4.5]_*$ stellar images is the same as that which we see in the optical. There are no hidden bars, and bars that are present look the same in the infrared as in the optical (DDO 154, NGC 3738, NGC 4214, NGC 4449, NGC 6822). The stellar populations that are emphasized may be different in different passbands, but the structure of the galaxy is the same for different stellar populations in Im galaxies. One exception is IC 10 that lies at low Galactic latitude; a Galactic dust lane obscures the western part of the galaxy in the optical and the stars both in IC 10 and in the Milky Way behind the dust lane become apparent in the $3.6 \mu\text{m}$

image (see Figure 3). Another exception is NGC 3738. This Im galaxy with intense central star formation looks similar in V and $[3.6]$, but is less round in J at outer isophotes (Hunter & Elmegreen 2006).

3.3. Embedded sources

On a small scale there are some differences between the $[3.6]$ and V band that are apparent in 4 of the galaxies: IC 10, NGC 3738, NGC 4214, and NGC 4449. These four galaxies are distinguished from the rest of the sample by intense star formation; and in three of these systems star formation is nearly galaxy-wide (in NGC 3738 the star formation is concentrated to the center of the galaxy). These galaxies are shown in Figure 3 where the $3.6\ \mu\text{m}$ and V images have been combined in false color and regions of $3.6\ \mu\text{m}$ excess have been identified in different passbands.

In IC 10 the main source of excess $3.6\ \mu\text{m}$ emission is along the southeast edge of the highest surface brightness part of the galaxy. In the optical there is an obvious dust lane running northeast to southwest there, and we see enhanced diffuse $3.6\ \mu\text{m}$ emission from this region (numbered D1 in Figure 3). Along the northwest edge of this dust lane there is a region of star formation, prominent in $\text{H}\alpha$. In the $3.6\ \mu\text{m}$ image there are 3 knots, all resolved with respect to a point source. Two of these knots are visible in V (numbered 1 and 3 in Figure 3); the third is not seen in V (numbered 2 in Figure 3). On the southeast edge of the dust lane, there is an unresolved point source apparent at $3.6\ \mu\text{m}$ and not in V (numbered 4 in Figure 3).

In NGC 3738, we see a source that is bright in $3.6\ \mu\text{m}$ in the northern part of the central star-forming complex (numbered 1 in Figure 3). This source consists of two resolved knots. The western of the two knots is associated with bright $\text{H}\alpha$ emission and faint V -band emission. The eastern of the two knots has fainter $\text{H}\alpha$ emission and is not visible in V .

In NGC 4214 the most interesting excess $3.6\ \mu\text{m}$ emission is associated with an arc of star forming regions in the southern part of the galaxy. Region 1 consists of two knots and region 2 is one knot, all resolved with respect to a point source. They are seen in both $3.6\ \mu\text{m}$ and V , but there appears to be excess $3.6\ \mu\text{m}$ emission in and around these knots. Region D1 is a diffuse extension southward of knot 1 seen only in $3.6\ \mu\text{m}$. In addition to this star forming region, diffuse excess $3.6\ \mu\text{m}$ emission is associated with a dust patch in the northwest part of the galaxy (numbered D2 in Figure 3). We note that a similar dust patch obvious in the optical located at $12^{\text{h}}\ 15^{\text{m}}\ 39^{\text{s}}.3$, $36^{\circ}\ 19'\ 22''$ (J2000.0) does not show excess $3.6\ \mu\text{m}$ emission.

NGC 4449 contains numerous regions of excess $3.6\ \mu\text{m}$ emission. Some of this emission is filamentary and suggests a nebular origin. The similarity between the $3.6\ \mu\text{m}$ image and the non-star-subtracted $4.5\ \mu\text{m}$ image suggests that some of the emission at $3.6\ \mu\text{m}$ could be from hot dust or the $3.3\ \mu\text{m}$ PAH feature. Several regions of particular note are circled in Figure 3. Region 1, located near the galaxy center, is a resolved knot in $3.6\ \mu\text{m}$ and a dust patch in the optical. Regions D1 and D2 are diffuse $3.6\ \mu\text{m}$ emission regions associated with optical dust patches. Region D3, also diffuse, lies along the outer edge of a large star forming complex in the northern part of the galaxy.

We have measured the brightnesses of the regions identified in Figure 3 in UBV , $3.6\ \mu\text{m}$, and $4.5\ \mu\text{m}$, where possible, and the photometry is given in Table 4. Numbered regions are resolved knots in $3.6\ \mu\text{m}$, and are photometered using circular apertures with background subtraction determined from an annulus just beyond the aperture. The radius of the aperture is given in Table 4. Regions designated with a “D” in the table are diffuse regions that were photometered using a polygon whose shape was guided by the $V/3.6\ \mu\text{m}$ image. For the diffuse regions, there was no background subtraction. The UBV photometry was calibrated using the data of Hunter & Elmegreen (2006); the IRAC photometry was calibrated as discussed in §2. In Table 4 we also give the $3.6\ \mu\text{m}$ surface brightness of each region in the aperture. Since the calibration of the $[3.6]$ image is relative to Vega, we report the surface brightness in terms of the luminosity of Vega at $3.6\ \mu\text{m}$ per square parsec, using a distance to Vega of 7.6 pc. When the output of star clusters or composite stellar populations in the $3.6\ \mu\text{m}$ passband are better known, one can relate these surface brightnesses to a more appropriate stellar population.

From Table 4 one can see that most of the $[3.6] - [4.5]_*$ colors are those of early K stars, similar to the integrated colors of Im galaxies. The resolved knots have surface brightnesses comparable to 30–500 stars like Vega per square parsec, while the diffuse regions have surface brightnesses of 50–120 Vegas per square parsec. Photometry of star forming complexes in 4 galaxies suggest that a $V - [3.6]$ color of 2.3 ± 0.5 is typical of young regions. Given that, the knots in IC 10 have total A_V of roughly 4.4–6.8 mag, and reddening-corrected M_V of order -9.5 to -9.9 . Thus, they are potentially populous star clusters (Billett et al. 2002), although the reddening corrected UBV colors do not make very much sense compared to cluster evolutionary tracks. The two knots in NGC 3738 have an A_V of order 2.7 if the intrinsic $V - [3.6]$ color is 2.3 mag. That would yield an M_V of -12.5 , potentially placing it in the realm of a super star cluster. Knots 1 and 2 in NGC 4214 would each have A_V of 0.5 mag, and M_V of -11.5 and -11.9 . Thus, they could also be massive star clusters. The bright diffuse regions in NGC 4449 are probably patches of stellar disk.

The presence of embedded luminous clusters in four of our galaxies is consistent with

the star formation seen in other tracers. The fact that there are only a few of these clusters in small galaxies with overall star formation rates of $\sim 0.05 \text{ M}_\odot \text{ yr}^{-1} \text{ kpc}^{-1}$ is not surprising. In IC 10, the SFR is $0.049 \text{ M}_\odot \text{ yr}^{-1} \text{ kpc}^{-1}$ normalized to the exponential disk radius of $R_D^V = 1.27'$ or 0.37 kpc . This gives a total star formation rate of $0.02 \text{ M}_\odot \text{ yr}^{-1}$. If, in an extreme case, all of this star formation were in the form of populous clusters, each having a mass of $10^4 \text{ M}_\odot$, then to get one such cluster still in the embedded phase requires an obscuring time of $10^4/0.02 = 5 \times 10^5 \text{ yr}$. In fact, $10^4 \text{ M}_\odot$ clusters are relatively rare. For the usual $M^{-2}dM$ cluster mass function, the fraction of the total star formation rate from a decade interval of cluster mass is $\log(10)/\log(M_U/M_L) = 1/\log(M_U/M_L)$ for upper and lower cluster mass limits M_U and M_L . Typically, $\log(M_U/M_L) \sim 4$ to 5 , depending on how a cluster is defined. Thus $\sim 20\%$ of the star formation rate can be in “populous” clusters like the embedded one in IC 10. This means the obscuration time is $10^4/(0.02 * 0.2) = 2.5 \times 10^6 \text{ yr}$. This is actually a reasonable timescale considering observations of local obscured clusters (e.g. Lada & Lada 2003). The total star formation rates in NGC 3738 and NGC 4214 are $0.09 \text{ M}_\odot \text{ yr}^{-1}$ and $0.13 \text{ M}_\odot \text{ yr}^{-1}$, significantly larger than in IC 10, so it is not unreasonable that the largest embedded clusters are also more massive than in IC 10, if the obscuration time is about the same. This follows from statistically sampling a universal cluster mass function. All of the other Im galaxies without bright embedded clusters could have their most massive clusters at later phases in their cloud break out, or they could have lower total star formation rates.

3.4. Surface photometry

The optical radial light profiles of Im galaxies are those of exponential disks (Hodge 1971), and we see this same form in the mid-IR. The $[3.6]$ or $[4.5]_*$ azimuthally averaged stellar surface photometry is shown for each galaxy in Figure 1. The central surface brightnesses μ_0 and disk scale lengths R_D derived from exponential disk fits to the surface photometry are given in Tables 2 and 3, and are plotted against the V -band measurements in Figures 4 and 5. The central surface brightnesses and disk scale lengths measured from the IRAC images are clearly correlated with those measured from V -band images. That the disk scale lengths are similar measured at different wavelengths implies that the structure of the disk is usually the same for old and low mass stars as for younger and more massive stars.

We have previously found complex V -band surface brightness profiles in approximately one-quarter of our survey of Im and Sm galaxies (Hunter & Elmegreen 2006). These profiles are double exponentials: one exponential fits the surface brightnesses in the central regions of the galaxy disk and an exponential with a different slope fits the outer parts. The most

common type of profile is one in which the outer part of the galaxy exhibits an exponential form that is steeper than that in the central part of the galaxy. In a few systems the inner profile was actually flat, and in one system the outer profile was shallower than the one at smaller radii. In our IRAC sub-sample, we have 4 galaxies that exhibit one of these complex profiles: DDO 165, IC 10, NGC 3738, and VIIZw 403. In Figure 1 one can see that the profiles in $[3.6]$ are very similar to those in V band: The breaks in the profiles are obvious and occur at, or nearly at, the same radii, and the slopes in the profiles before and after the breaks are similar. Therefore, the breaks and changes in the radial surface density of stars is the same in different stellar populations within the galaxy.

Azimuthally-averaged colors are also shown as a function of radius for each galaxy in Figure 1. We find that 68% (13 of 19) of the Im galaxies have $V - [3.6]$ or $V - [4.5]_*$ colors that are constant with radius. Thus, the mixture of young and old stars is the same throughout the galaxy disks. Both BCDs with these colors show a gradient, at least in the galaxy center, getting redder with distance from the center of the galaxy. This is consistent with the central concentration of intense star formation in these systems. For $[3.6] - [4.5]_*$, the color is constant with radius in 85% (11 of 13) of the Im galaxies and all 3 of the BCDs.

4. Hot Dust

4.1. Spatial distribution

The star-subtracted $[4.5]_{\text{dust}}$ image shows the distribution and brightness of hot dust emission in a galaxy. The distribution of the hot dust in our sample of galaxies is shown as contours superposed on $H\alpha$ images in Figure 1. We note that the hot dust is always much less extensive than the PAHs as seen in $8.0 \mu\text{m}$ -band emission. In fact, most, and probably all, regions of hot dust are associated with bright H II regions. There are a few regions in several galaxies that are not associated with H II regions, but these are most likely foreground or background objects and not associated with the object galaxy at all. H II regions are where the concentrations of UV photons and dust are located, so this implies that young, massive stars are the source of heating of this component of the dust.

We have measured the $H\alpha$ surface brightness limit at which hot dust is detected $\Sigma_{H\alpha}^{[4.5]_{\text{dust}}}$. The average is $8 \times 10^{29} \text{ erg s}^{-1} \text{ pc}^{-2}$ for the Im galaxies with a factor of 8 variation among galaxies. The average for 2 BCDs is $11 \times 10^{29} \text{ erg s}^{-1} \text{ pc}^{-2}$, with a factor of 1.4 variation. The $H\alpha$ surface brightness limits are plotted as a function of oxygen abundance and area-normalized star formation rate in the bottom panels of Figure 6. There is no correlation with oxygen abundance, but there is a clear correlation with star formation rate: as the star

formation rate goes down, the $H\alpha$ surface brightness limit at which hot dust is detected also goes down. Thus, there is no single surface brightness that predicts where hot dust will be found for all Im galaxies.

4.2. Integrated properties

We have detected $[4.5]_{\text{dust}}$ emission in 69% (11 of 16) of the galaxies with observations in both $3.6\ \mu\text{m}$ and $4.5\ \mu\text{m}$ passbands. The integrated brightness of the hot dust emission in the $4.5\ \mu\text{m}$ passband is given as an absolute magnitude in Table 2. The oxygen abundance, area-normalized star formation rate, and integrated $(U-B)_0$ color are plotted as a function of this absolute magnitude $M_{[4.5]_{\text{dust}}}$ in Figure 7. There is a slight increase in the amount of hot dust with increasing star formation rate (correlation coefficient 0.6). This is consistent with the observation discussed in the previous section that hot dust in Im galaxies is primarily detected in H II regions above a particular surface brightness level. There is no correlation of $M_{[4.5]_{\text{dust}}}$ with $(U-B)_0$. There is a slight correlation with oxygen abundance (correlation coefficient 0.5), probably reflecting the luminosity-metallicity relationship seen for galaxies since there is a correlation between $M_{[4.5]_{\text{dust}}}$ and the stellar absolute magnitude $M_{[3.6]}$.

5. PAHs

5.1. Integrated properties

The star-subtracted $[8.0]_{\text{PAH}}$ passband image contains emission from PAHs, as well as from warm dust (Rosenberg et al. 2006). We have detected this emission in 73% (16 of 22) of the galaxies in our sample. The integrated brightness in the non-stellar $[8.0]_{\text{PAH}}$ image converted to an absolute magnitude $M_{[8.0]_{\text{PAH}}}$ is given in Tables 2 and 3 and plotted against integrated M_V , several colors, the area-normalized star formation rate, and oxygen abundance in Figure 8. PAH emission increases with increasing V -band (or $3.6\ \mu\text{m}$) luminosity. There is also a correlation between $M_{[8.0]_{\text{PAH}}}$ and $[5.8]_{\text{PAH}}-[8.0]_{\text{PAH}}$ color in the sense that, as the PAH emission in the $[8.0]_{\text{PAH}}$ image increases, it does so faster than the emission in $[5.8]_{\text{PAH}}$ increases. No $[5.8]_{\text{PAH}}$ emission is detected in galaxies with $M_{[8.0]_{\text{PAH}}}$ below -17 . This represents a detection limit for $5.8\ \mu\text{m}$ non-stellar emission in our galaxy sample.

In Figure 9 we plot the color $[3.6]-[8.0]_{\text{PAH}}$ against the $8.0\ \mu\text{m}$ PAH brightness, area-normalized star formation, and oxygen abundance. The color $[3.6]-[8.0]_{\text{PAH}}$ is a measure of the $[8.0]_{\text{PAH}}$ PAH emission relative to the mid-IR stellar emission. Most of the galaxies in our sample have $[3.6]-[8.0]_{\text{PAH}}$ colors in the range 0–3 mag. Hogg et al. (2005) found that

star-forming galaxies have larger PAH emission relative to stars, but that low luminosity systems have unusually low PAH-to-star ratios. In a study of galaxies spanning the Hubble sequence, Pahre et al. (2004) found that later type galaxies have higher PAH emission relative to starlight. They measure $[3.6] - [8.0]_{\text{PAH}} \sim 2$ for types greater than $T=3$. In the disk of M81, Willner et al. (2004) found a relatively constant $[3.6] - [8.0]_{\text{PAH}}$ color with radius at a value of about 0.5. The high star formation rate dwarf sample of Rosenberg et al. (2006) has colors 1–3 mag. Thus, our sample covers the range of colors seen in other samples of Im or low luminosity galaxies.

There is a correlation between PAH emission and star formation rate, seen in both Figure 8 and Figure 9: PAH emission increases relative to stellar emission as the star formation rate per unit area increases. Roussel et al. (2001) and Förster Schreiber et al. (2004) found a correlation between star formation rate and the 5–8.5 μm emission in spiral and starburst systems, and Dale et al. (2005) have suggested that the $[8.0]_{\text{PAH}}$ PAH emission generally traces the star formation rate with factors of 10–20 variations in galaxies. Rosenberg et al. (2006) also saw this correlation in their sample of starburst dwarfs.

The correlation of PAH emission with star formation rate suggests that the stars that ionize the gas also heat the PAHs. This is consistent with the spatial distribution discussed in §5.3 in which, for most galaxies, the PAH emission is found in the brightest H II regions. Although cool stars can heat PAHs too (Uchida et al. 1998, Li & Draine 2002), in these galaxies the primary source of heating are the concentrations of young, massive stars in H II regions. The exceptions in our sample are the galaxies with very intense, galaxy-wide star formation. In those systems the PAH emission is much more extended. But, the wide-spread nature of the star formation in these systems, coupled with the porosity that comes from concentrations of massive stars blowing holes in the ISM, allows UV photons from the star-forming regions to escape to large distances. Thus, the heating source for the PAHs is likely the same in all of the galaxies in our sample. However, the strong UV fluxes can also destroy or ionize PAHs (Boulange et al. 1988, Vestraete et al. 1996, Tacconi-Garmon et al. 2005). Therefore, we would expect that very high spatial resolution imaging would reveal that the PAH emission is actually located in the outer parts of the H II regions, as is customarily found (for example, Churchwell et al. 2004, Marston et al. 2004, Rho et al. 2006).

However, Spoon (2003) has suggested that on galactic scales the PAH emission better traces the B stars rather than sites of younger star formation. In a study of mid-IR emission in M51, Calzetti et al. (2005) state that the UV photons that heat the PAHs come from recent past star formation (≤ 100 Myr old), as well as current star formation (≤ 10 Myr). Therefore, the $[8.0]_{\text{PAH}}$ PAH emission is correlated with star formation but not directly proportional to the number of ionizing photons. For our sample, in Figure 8 we see that the $[8.0]_{\text{PAH}}$

PAH brightness is independent of integrated galactic $(U-B)_0$. $(U-B)_0$ is dominated by the young stellar component of the galaxy, but since it contains a contribution from relatively older, redder B stars as well as OB stars in star-forming regions, we might expect $(U-B)_0$ to become bluer (up to the $(U-B)_0$ color of an O star) as PAH emission increases if the only contribution to the PAH heating were from OB stars in H II regions. That we do not see such a trend suggests that B stars outside of H II regions are also contributing to PAH heating in Im galaxies. On the other hand, in Figure 9 galaxies with a greater proportion of PAH emission relative to stellar $3.6\ \mu\text{m}$ emission are in fact slightly bluer in $(U-B)_0$. (There is no separate correlation between $M_{[3.6]}$ or $M_{[8.0]_{\text{PAH}}}$ and $(U-B)_0$; see Figures 2 and 8). However, the correlation is shallow. All of this together suggests that young, massive stars in star-forming regions are a major source of heating of PAHs in Im galaxies, but that B stars outside of star-forming regions also contribute.

Engelbracht et al. (2005) has found that the $[8.0]_{\text{PAH}}$ PAH emission decreases relative to the $24\ \mu\text{m}$ dust emission for galaxies with metallicity less than $1/3$ – $1/5$ solar ($12 + \log \text{O}/\text{H} \leq 8.2$). Thus, we expect most, if not all, of the galaxies in our sample to have depressed PAH emission, and as a group this may be true. However, we have explored the relationship between metallicity and PAH emission within this low metallicity sample. In Figures 8 and 9 we see that galaxies with higher oxygen abundances have higher $[8.0]_{\text{PAH}}$ PAH emission and more PAH emission relative to that of stars at $3.6\ \mu\text{m}$. Rosenberg et al. (2006) also see a correlation between $M_{[8.0]_{\text{PAH}}}$ and oxygen abundance for their star-forming dwarfs, and the slopes of our relationships are very similar: We measure 0.08 ± 0.02 , and they measure 0.07 . Rosenberg et al. and Hogg et al. (2005) also see a slight trend with a lot of scatter in a plot of $[3.6] - [8.0]_{\text{PAH}}$ with oxygen abundance for low luminosity systems.

5.2. Surface photometry

In Figure 1 we explore the variation of non-stellar $[5.8]_{\text{PAH}} - [8.0]_{\text{PAH}}$ color with radius through azimuthally averaged surface photometry for each galaxy. We see that in 67% (4 of 6) of the Im galaxies and one of two BCDs this color is constant with radius. In a study of the disk of M81, Willner et al. (2004) found that the $[5.8]_{\text{PAH}} - [8.0]_{\text{PAH}}$ color of non-stellar emission varies between about 0 and 0.5 with radius.

5.3. Two-dimensional spatial distribution

We illustrate the distribution of PAH emission relative to the star formation activity as seen in nebular $H\alpha$ emission in Figure 1. There we have superposed a low surface brightness contour of $[8.0]_{\text{PAH}}$ PAH emission on our $H\alpha$ image. We see that in most of the galaxies the PAH emission is associated with the brighter H II regions. This is also what we found from a study of a smaller sample of Im galaxies with *ISO* (Hunter et al. 2001). As discussed above, this implies that the young, massive stars are the primary heating source for the PAH and dust emission in the $[8.0]_{\text{PAH}}$ passband in Im galaxies. This is in contrast to the study of M81 by Willner et al. (2004) who found that $[8.0]_{\text{PAH}}$ PAH emission corresponds best with dust lanes in that spiral disk.

The average $H\alpha$ surface brightness limit for detecting PAH emission, excluding galaxies with extended emission, is $3 \times 10^{29} \text{ ergs s}^{-1} \text{ pc}^{-2}$ with a factor of 12 variation for the Im galaxies. For the two BCDs, the average limit is $10 \times 10^{29} \text{ ergs s}^{-1} \text{ pc}^{-2}$ with a factor of 10 variation. These limits are plotted as a function of oxygen abundance and area-normalized star formation rate in the top panels of Figure 6. Like the hot dust $H\alpha$ surface brightness limits, the lower the star formation rate, the lower the $H\alpha$ surface brightness limit at which PAH emission is detected. However, unlike that for hot dust emission, there is a correlation between the limits for PAH emission and oxygen abundance: Galaxies with higher metallicities have lower $H\alpha$ surface brightness limits for PAH emission.

Four galaxies in our sample have PAH emission that is much more extensive than location in a few discrete H II regions. These galaxies are IC 10, NGC 3738, NGC 4214, and NGC 4449. Not surprisingly, these galaxies are all characterized by intense, galaxy-wide star formation as well. In these galaxies the PAH emission is highly extended, even into the outer parts of the galaxy. NGC 6822 also has $8.0 \mu\text{m}$ emission outside of H II regions, although its emission is more localized and not as extensive as those in the other four galaxies.

Tacconi-Garman et al. (2005) found no correlation between star-forming regions and PAH emission on local scales in a study of two starburst systems. This is not the case in our sample where there is a fairly high degree of correlation between $H\alpha$ and PAH emission although the relative strengths vary. For four of the galaxies with extended PAH emission we have plotted azimuthally averaged radial profiles of the $H\alpha$ emission relative to the $[8.0]_{\text{PAH}}$ PAH emission in Figure 10. (Because of Milky Way cirrus, we do not have measurements in the $8.0 \mu\text{m}$ band for NGC 6822). We see that in NGC 3738, NGC 4214, and the inner part of NGC 4449 the ratio of $H\alpha$ to PAH emission remains relatively constant with radius. In the outer part of NGC 4449 and over most of IC 10 this ratio decreases with radius; the average PAH emission is stronger than we would expect for the $H\alpha$ emission at those radii.

We also see this on a local level. Figure 11 displays the $H\alpha$ and $[8.0]_{\text{PAH}}$ PAH images in false color, with $H\alpha$ in green and $[8.0]_{\text{PAH}}$ in red, for the 5 galaxies with extra- $H\text{ II}$ region PAH emission. One can see that most of the PAH emission is correlated with $H\alpha$ emission, but not always. There are places within the galaxies where PAH emission is found without $H\alpha$ or the $H\alpha$ brightness is very low. The center of NGC 6822 is an example of high PAH emission with almost no $H\alpha$ emission. The NGC 6822 field is contaminated by Galactic cirrus, but the region in question is higher in surface brightness and smaller in size than the large-scale pattern one sees across the image. Spoon (2003) and Calzetti et al. (2005) have suggested that B stars no longer associated with $H\text{ II}$ regions can be a major source of heating of the PAHs in galaxies. This could be the case for this region in NGC 6822, because that region of the disk is slightly redder than its surroundings. The lack of star formation means there are few dense molecular clouds. There can still be PAH emission, however, if there are diffuse clouds irradiated by local B stars and more remote OB stars. The inner part of NGC 6822 could therefore be unusual in the sense that most of the ISM is in a diffuse form that does not readily form stars. The same could be true for IC 10 where the extended regions of high PAH-to- $H\alpha$ emission could be irradiated by local B stars and distant OB stars. This is somewhat easier to understand in IC 10 than in NGC 6822 because IC 10 has many gas holes through which UV radiation can travel long distances (Wilcots & Miller 1998).

5.4. PAH emission in shells and supershells

Figure 12 illustrates the fact that PAH emission is also found along the rims of gas shells and filaments. One can see this in IC 10 where the two gas shells to the northwest of the highest surface brightness part of the galaxy ($H\alpha$ shells numbered 5 and 7 by Wilcots & Miller 1998; see their Figure 11) have PAH emission mixed with $H\alpha$ emission. In fact these shells are more complete in the $[8.0]_{\text{PAH}}$ image than in $H\alpha$. This is especially true of shell number 5 where the $H\alpha$ emission is concentrated in $H\text{ II}$ regions embedded in the $H\text{ I}$ shell, while the $[8.0]_{\text{PAH}}$ image shows fainter extra- $H\text{ II}$ region emission along the shell. These shells, seen in $H\text{ I}$ as well, are about 200-290 pc in diameter (for a distance of 1 Mpc; Wilcots & Miller 1998) and represent the mechanical energy input of concentrations of massive stars to their surroundings. Cannon et al. (2005) also see PAH emission associated with a gas shell in the Im galaxy IC 2574.

An even more extreme example is found in NGC 4449. There one can see PAH emission associated with a supergiant gas filament in the northwest part of the outer galaxy. These are filaments 5 and 6 of Hunter & Gallagher (1992; and best seen in Figures 1 and 4 of Hunter & Gallagher 1997). The $H\alpha$ filaments form a coherent structure 1.7 kpc long, and

filament 6 falls along the inner edge of a 1.8 kpc diameter H I supershell. PAH emission is found in broken patches along these filaments. At the base of this supershell near the center of the galaxy ionized gas is expanding at 30 km s^{-1} . These and other structures create a picture in which intense star-forming events have created multiple supergiant shells in the ISM of NGC 4449 (Hunter & Gallagher 1997). Thus, PAHs are not destroyed in the shell formation process, but rather are swept up into shells and filaments along with the gas. Tacconi-Garman et al. (2005) suggest that this means that galaxies are capable of enriching their haloes and the intergalactic medium with dust.

The PAH emission is not always spatially coincident with the H I or H α emission in the shells. In IC 10's shell number 5 and the northwest edge of shell number 7 (see Figure 12 of Wilcots & Miller 1998) the H I, H α , and PAH emission are spatially coincident. However, along the northern edge of shell 7, the H α emission is located along the inner edge of the H I ridge while the $8.0 \mu\text{m}$ emission extends further into the ridge by 10–30 pc. The $8.0 \mu\text{m}$ appears brightest where the H α is brightest in this shell. Also, in the far northeast, the H I emission continues for a distance of at least twice the field of view in Figure 11 with a column density of $\sim 10^{21} \text{ cm}^{-2}$ (Wilcots & Miller 1998), but there is little $8.0 \mu\text{m}$ emission in the northeastern part of our field of view. On the other hand, in the western part of Figure 11, there is a lot of dispersed $8.0 \mu\text{m}$ covering a broad area where the H I is lacking.

In NGC 4449 the H α and PAH emission are spatially coincident in filament 5, to within our ability to measure the positions (about $2''=38 \text{ pc}$), but both lie along the inner edge of the H I ridge, offset about 265 pc ($14''$) from the ridge center. (The FWHM of the H I beam is $15''$). Along the H I supershell, the diffuse H α and PAH emission along the northwest edge appear to be spatially coincident, although the center of such faint, diffuse emission is difficult to accurately determine. The center of the H I ridge is offset further out by 190 pc or $0.3R_{\text{shell}}$. A different situation pertains along the southwest (bottom) edge of the H I supershell: The H I and $[8.0]_{\text{PAH}}$ emission are spatially coincident while the H α lies along the inner H I edge, offset by 110–210 pc ($0.15\text{--}0.3R_{\text{shell}}$) from the center of the ridge.

These differences between $8 \mu\text{m}$ and H I could be from a combination of several effects. First, the $8.0 \mu\text{m}$ emission from PAHs requires both gas and starlight, so there can be a decrease in $8.0 \mu\text{m}$ relative to H I where the radiation field decreases in the presence of neutral gas. For example, in IC 10, the decrease in $8.0 \mu\text{m}$ compared to H I in the northern part of shell 7 may occur because of extinction inside the ridge. Similarly, the relative decrease of $8.0 \mu\text{m}$ in the northeast of IC 10 may occur because of the great distance from the radiation source, which is concentrated in the center of the galaxy. The concentration of $8.0 \mu\text{m}$ to the irradiated parts of H I shells in NGC 4449 is probably a similar effect. A second process that could change the relative emission strengths of $8.0 \mu\text{m}$ and H I is diffuse

ionization. The extended diffuse 8.0 μm to the west of IC 10 (see above) corresponds to a giant H I hole that could be mostly diffuse ionized gas; there is a bright Wolf-Rayet star inside this H I hole (Massey et al. 1992; Wilcots & Miller 1998). The PAHs would then be dispersed and irradiated in the ionized gas. These two effects should cause a different relative abundance of ionized versus neutral PAH emission, which might be detectable in IR spectra.

6. Summary

We have examined mid-IR images of 22 irregular galaxies obtained from the *Spitzer* archives. The galaxies are representative of normal systems of the Im, BCD, and Sm classes and span the range of luminosities and star formation rates found in larger samples. From the [3.6] and [4.5] images we extract information about the structure of the stellar disk and its stellar populations. The star-subtracted [4.5]_{dust} image is used to examine the distribution and integrated emission from hot dust. The nebular [5.8]_{PAH} and [8.0]_{PAH} images are used to explore the properties of PAH and hot dust emission. The two-dimensional distributions, integrated magnitudes and colors, and azimuthally averaged surface photometry are compared to optical images and measurements, including H α images that trace the current star formation activity and location of luminous, hot stars.

From the stellar images, we find that generally there are no hidden bar structures in IRAC bands. Four of the Im galaxies have optically-hidden bright IR sources, with M_V at least as bright as -9.5 to -12.5 mag, placing them in the range for populous and super star clusters. Abrupt changes in some radial stellar density profiles seen in the optical, suggesting double-exponential disks, are also seen in the near-IR. For most Im and all BCD and Sm galaxies, the overall structure of the galaxy is the same in the different passbands. Furthermore, the colors suggest that in most Im galaxies the mixture of young and old stars is the same throughout the galaxy disk.

The hot dust seen in the star-subtracted [4.5]_{dust} image is detected in 2/3 of the galaxies. The hot dust is found only in the higher surface brightness H II regions, where the concentrations of UV photons as well as dust are located. This implies that young, massive stars are the source of heating of this component of the dust in Im and BCD galaxies. Furthermore, galaxies with higher area-normalized star formation rates have more dust emission.

PAH plus dust emission in the star-subtracted [8.0]_{PAH} image is detected in nearly 3/4 of our sample of galaxies. As the area-normalized star formation rate increases, the PAH emission increases relative to the starlight. Furthermore, except for a few galaxies, the PAH emission is found only in the brightest H II regions. The exceptions are galaxies with intense,

wide-spread star formation, and in those systems the PAH emission is much more extended, even to the outer parts of the galaxy. The correlation with star formation activity suggests that, on average, the concentrations of young, massive stars that ionize the gas also primarily heat the PAHs in Im-type galaxies. However, on a local level, the ratio of PAH emission to $H\alpha$ emission is not constant.

We see PAH emission associated with the rims of gas shells and super-giant filaments that have resulted from the intense mechanical energy input from concentrations of massive stars. This implies that PAHs are not destroyed in the shell formation process, but are swept up into the shells and filaments along with the gas even to the outer parts of a galaxy.

E. M. would like to thank Kathy Eastwood and the 2005 Research Experience for Undergraduates program at Northern Arizona University which is funded by the National Science Foundation (NSF) under grant AST-0453611. We appreciate educational exchanges about IRAC with M. Pahre, S. Willner, and M. Ashby. We thank M. Fitzpatrick for help with an IRAF problem. Funding for this work was provided to D. A. H. by the Lowell Research Fund and the NSF through grant AST 02-04922. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center (IPAC)/California Institute of Technology, funded by the National Aeronautics and Space Administration (NASA) and NSF. This research has also made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with NASA.

Facilities: Spitzer Space Telescope, Lowell Observatory.

REFERENCES

- Allamandola, L. J., Tielens, G. G. M., & Barker, J. R. 1989, *ApJS*, 71, 733
- Billett, O. H., Hunter, D. A., & Elmegreen, B. G. 2002, *AJ*, 123, 1454
- Boselli, A., Lequeux, J., Sauvage, M., Boulade, O., Boulanger, F., Cesarsky, D., Dupraz, C., Madden, S., Viallefond, F., & Vigroux, L. 1998, *A&A*, 335, 53
- Boulanger, F., Beichman, C., Desert, F. X., Helou, G., Perault, M., & Ryter, C. 1988, *ApJ*, 332, 328
- Calzetti, D., et al. 2005, *ApJ*, 633, 871

- Cannon, J. M., et al. 2005, *ApJ*, 630, L37
- Churchwell, E., et al. 2004, *ApJS*, 154, 322
- Dale, D. A., et al. 2005, *ApJ*, 633, 857
- de Blok, W. J. G., & Walter, F. 2000, *ApJ*, 537, L95
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H., Buta, R., Paturel, G., & Fouqué, P. 1991, *Third Reference Catalogue of Bright Galaxies* (New York, Springer-Verlag)
- Draine, B. T. 2005, in *The Spitzer Science Center 2005 Conference: Infrared Diagnostics of Galaxy Evolution*, in preparation
- Engelbracht, C. W., Gordon, K. D., Rieke, G. H., Werner, M. W., Dale, D. A., & Latter, W. B. 2005, *ApJ*, 628, L29
- Fazio, G. G., et al. 2004, *ApJS*, 154, 10
- Förster Schreiber, N. M., Roussel, H., Sauvage, M., & Charmandaris, V. 2004, *A&A*, 419, 501
- Galliano, F., Madden, S. C., Jones, A. P., Wilson, C. D., Bernard, J.-P., & Le Peintre, F. 2003, *A&A*, 407, 159
- Hodge, P. W. 1971, *ARA&A*, 9, 35
- Hogg, D. W., et al. 2005, *ApJ*, 624, 162
- Houck, J. R., et al. 2004, *ApJS*, 154, 211
- Hunter, D. A., & Elmegreen, B. G. 2004, *AJ*, 128, 2170
- Hunter, D. A., & Elmegreen, B. G. 2006, *ApJS*, 162, 49
- Hunter, D. A., et al. 2001, *ApJ*, 553, 121
- Hunter, D. A., & Gallagher, J. S., III 1992, *ApJ*, 391, L9
- Hunter, D. A., & Gallagher, J. S., III 1997, 475, 65
- Hunter, D. A., Gallagher, J. S., III, Rice, W. L., & Gillett, F. C. 1989, *ApJ*, 336, 152
- Hunter, D. A., Gillett, F., Gallagher, J. S., III, Rice, W. L., and Low, F. 1986, *ApJ*, 303, 171

- Hunter, D. A., & Hoffman, L. 1999, *AJ*, 117, 2789
- Hunter, D. A., van Woerden, H., & Gallagher, J. S. 1999, *AJ*, 118, 2184
- Kennicutt, R. C., Jr., et al. 2003, *PASP*, 115, 928
- Kennicutt, R. C., & Skillman, E. D. 2001, *AJ*, 121, 14614
- Kniazev, A. Y., et al. 2000, *A&A*, 357, 101
- Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41 57
- Lee, H., & Skillman, E. D. 2004, *ApJ*, 614, 698
- Léger, A., & Puget, J. L. 1984, *A&A*, 137, L5
- Li, A., & Draine, B. T. 2002, *ApJ*, 572, 232
- Lu, N., et al. 2003, *ApJ*, 588, 199
- Madden, S. C. 2000, *New Astr Rev*, 44, 249
- Marston, A. P., et al. 2004, *ApJS*, 154, 333
- Massey, P., Armandroff, T. E., & Conti, P. S. 1992, *AJ*, 103, 1159
- Mo, H. J., Mao, S., & White, S. D. M. 1998, *MNRAS*, 295, 319
- Nikolaev, S., & Weinberg, M. D. 2000, *ApJ*, 542, 804
- Pahre, M. A., Ashby, M. L. N., Fazio, G. G., & Willner, S. P. 2004, *ApJS*, 154, 235
- Pilyugin, L. S. 2001, *A&A*, 374, 412
- Reach, W. T., et al. 2005, *PASP*, 117, 978
- Regan, M. W., et al. 2004, *ApJS*, 154, 204
- Rho, J., Reach, W. T., Lefloch, B., & Fazio, G. 2006, *ApJ*, in press
- Richer, M. G., & McCall, M. L. 1995, *ApJ*, 445, 642
- Rosenberg, J. L., Ashby, M. L. N., Salzer, J. J., & Huang, J.-S. 2006, *ApJ*, 636, 742
- Roussel, H., Gauvage, M., Vigroux, L., & Bosma, A. 2001, *A&A*, 372, 427
- Skillman, E. D., Bomans, D. J., & Kobulnicky, H. A. 1997, *ApJ*, 474, 205

- Spoon, H. W. W. 2003, Ph.D. Thesis, Rijksuniversiteit Groningen
- Sturm, E., Lutz, D., Tran, D., Feuchtgruber, H., Genzel, R., Kunze, D., Moorwood, A. F. M., & Thornley, M. D. 2000, *A&A*, 358, 481
- Tacconi-Garman, L. E., et al. 2005, *A&A*, 432, 91
- Talent, D. 1980, Ph.D. Thesis, Rice University
- Thuan, T. X., Sauvage, M., & Madden, S. 1999, *ApJ*, 516, 783
- Uchida, K. I., Sellgren, K., & Werner, M. 1998, *ApJ*, 493, 109
- Verstraete, L., Puget, J. L., Falgarone, E., Drapatz, S., Wright, C. M., & Timmermann, R. 1996, *A&A*, 315, L337
- Wilcots, E. M., & Miller, B. W. 1998, *AJ*, 116, 2363
- Willner, S. P., et al. 2004, *ApJS*, 154, 222

Fig. 1.— Images and surface photometry for all of the galaxies in this study, two rows per galaxy and two galaxies per page. The images all show the same field of view for each galaxy. *Top left:* The geometrically transformed [3.6] or [4.5] image or the logarithm of the image. The contour is of an outer V -band isophote (Hunter & Elmegreen 2006). *Top middle:* An $H\alpha$ image (Hunter & Elmegreen 2004), if the galaxy has $H\alpha$ emission, is shown with a contour of [8.0]_{PAH} PAH emission (with stars removed as discussed in §2). If the galaxy has no detectable PAH emission, the contour is the same V -band contour shown in the upper left image. *Bottom left:* An $H\alpha$ image, if the galaxy has $H\alpha$ emission, is shown with a contour of [4.5]_{dust} hot dust emission (stars removed as discussed in §2), if the galaxy has detectable hot dust emission. *Bottom middle:* The surface photometry in V , $H\alpha$, [3.6] or [4.5]_{*}, and [8.0]_{PAH} PAH emission, if detected, as a function of radius is plotted here. The scale on the left vertical axis is for V band, and the scale on the right vertical axis is for $H\alpha$. The zero points of the IRAC surface photometry have been adjusted so that plots of the different quantities are easily distinguished, but the scales have all been set so that they cover the same logarithmic interval. The solid lines are fits to the exponential disks in V , J if observed, and [3.6] or [4.5]_{*}. *Right:* Plot of the colors as a function of radius from the azimuthally-averaged annular photometry. [The first of 11 parts is shown here.]

Fig. 2.— Integrated absolute V magnitude and several colors plotted as a function of absolute $[3.6]$ magnitude. The dashed lines are fits to the data: *Top panel:* $M_V = (-0.83 \pm 0.63) + (0.83 \pm 0.04)M_{[3.6]}$; *Second panel:* $(B-V)_0 = (-0.08 \pm 0.19) - (0.02 \pm 0.01)M_{[3.6]}$; *Third panel:* $(U-B)_0 = -0.31 \pm 0.16$; *Fourth panel:* $V-[3.6] = (-0.42 \pm 0.47) - (0.15 \pm 0.03)M_{[3.6]}$; *Bottom panel:* $[3.6] - [4.5]_* = -0.08 \pm 0.18$ (highly discrepant point is not included in average). $[4.5]_*$ is the stellar contribution to the $4.5 \mu\text{m}$ image; separation of the dust and stellar components is discussed in §2.

Fig. 3.— Comparison of $3.6\ \mu\text{m}$ images with other passbands for the 4 galaxies with regions of excess $3.6\ \mu\text{m}$ emission: IC 10, NGC 3738, NGC 4214, and NGC 4449. There are 5 panels for each galaxy. The 4 panels in black and white show the same field of view; the color panel generally shows a somewhat larger field of view. *Top right:* False color images comparing V (*green*) and $3.6\ \mu\text{m}$ (*red*). Regions of interest are circled. *Top left:* $V/3.6\ \mu\text{m}$ image. Photometered regions are circled and numbered as in Table 4, and the positions of the numbers and circles are retained in the other 3 black and white panels. White in this image represents excess $3.6\ \mu\text{m}$ emission. Some of the white regions in IC 10 and NGC 4214 are stars that were edited from the V -band image and not the $3.6\ \mu\text{m}$ image. *Top middle:* $\text{H}\alpha$ image from Hunter & Elmegreen (2004). The stellar continuum has been removed. *Bottom left:* $3.6\ \mu\text{m}$ image. Foreground stars and background objects have been removed from the NGC 3738 $3.6\ \mu\text{m}$ image. *Bottom middle:* V -band image. Foreground stars and background galaxies have been removed from the images except for IC 10.

Fig. 4.— Central surface brightnesses μ_0 from fits to the IRAC surface photometry compared to that measured in V band (Elmegreen & Hunter 2006). Galaxies with [3.6] images are shown in the upper panel, and those with only [4.5] images are shown in the bottom panel. The dashed lines are fits to the points: *Top panel:* $\mu_0^{[3.6]} = (-3.60 \pm 1.84) + (1.07 \pm 0.08)\mu_0^V$; *Bottom panel:* $\mu_0^{[4.5]*} = (-0.86 \pm 2.46) + (0.95 \pm 0.11)\mu_0^V$. [4.5]_{*} is the stellar contribution to the 4.5 μm image; separation of the dust and stellar components is discussed in §2. The BCD (HS 0822+3541) in the top panel has been left out of the fit.

Fig. 5.— Disk scale length measured from the IRAC [3.6] or [4.5]_{*} image $R_D^{[3.6]}$ or $R_D^{[4.5]_*}$ plotted against the disk scale length measured from the V -band surface photometry R_D^V (from Hunter & Elmegreen 2006). [4.5]_{*} is the stellar contribution to the 4.5 μm image; separation of the dust and stellar components is discussed in §2. For galaxies with double exponential profiles, one scale length is plotted as a solid symbol and the second one is plotted as an open symbol. The dashed line delineates a one-to-one correspondence and is not a fit to the data.

Fig. 6.— $H\alpha$ surface brightness limits at which hot dust measured in $[4.5]_{\text{dust}}$ images (*Bottom panels*) and $[8.0]_{\text{PAH}}$ PAH emission (*Top panels*) are detected in our sample galaxies are plotted against oxygen abundance and area-normalized star formation rate. $[4.5]_{\text{dust}}$ is the hot dust contribution to the $4.5 \mu\text{m}$ image; separation of the dust and stellar components is discussed in §2. The units of $\Sigma_{H\alpha}$ are $\text{ergs s}^{-1} \text{pc}^{-2}$. The oxygen abundance and star formation rates are integrated values over each galaxy. No limits were measured in galaxies with highly extended PAH emission. Dashed lines are fits to the data: *Top left*: $\log \Sigma_{H\alpha}^{[8.0]} = (41.78 \pm 2.52) - (1.55 \pm 0.32)(12 + \log \text{O}/\text{H})$; *Top right*: $\log \Sigma_{H\alpha}^{[8.0]} = (30.20 \pm 0.50) + (0.28 \pm 0.24)(\log \text{SFR}/\text{area})$; *Bottom left*: $\log \Sigma_{H\alpha}^{[4.5]_{\text{dust}}} = 29.99 \pm 0.25$; *Bottom right*: $\log \Sigma_{H\alpha}^{[4.5]_{\text{dust}}} = (30.37 \pm 0.16) + (0.26 \pm 0.08)(\log \text{SFR}/\text{area})$.

Fig. 7.— Integrated oxygen abundance, area-normalized star formation rate (Table 1), and $(U-B)_0$ plotted as a function of the integrated brightness of hot dust in the $[4.5]_{\text{dust}}$ images. $[4.5]_{\text{dust}}$ is the hot dust contribution to the $4.5 \mu\text{m}$ image; separation of the dust and stellar components is discussed in §2. The dashed lines are fits to the data: *Bottom panel:* $(U-B)_0 = -0.44 \pm 0.21$; *Middle panel:* $\log \text{SFR}/\text{area} = (-5.50 \pm 1.71) - (0.26 \pm 0.11)M_{[4.5]_{\text{dust}}}$; *Top panel:* $12 + \log \text{O}/\text{H} = (6.48 \pm 0.89) - (0.10 \pm 0.06)M_{[4.5]_{\text{dust}}}$.

Fig. 8.— Integrated M_V , PAH emission color $[5.8]_{\text{PAH}} - [8.0]_{\text{PAH}}$, optical color $(U - B)_0$, area-normalized star formation rate from Table 1, and oxygen abundance plotted as a function of the brightness of the PAH emission in the $[8.0]_{\text{PAH}}$ passband, reported as an absolute magnitude. Open symbols with arrows are upper limits. Dashed lines are fits to the data: *Top panel:* $M_V = (-4.59 \pm 0.97) + (0.59 \pm 0.05)M_{[8.0]_{\text{PAH}}}$; *Second panel:* $[5.8]_{\text{PAH}} - [8.0]_{\text{PAH}} = (-1.96 \pm 0.33) - (0.16 \pm 0.02)M_{[8.0]_{\text{PAH}}}$ (fit to only the Im galaxies); *Third panel:* $(U - B)_0 = -0.31 \pm 0.16$; *Fourth panel:* $\log \text{SFR/area} = (-6.96 \pm 1.70) - (0.27 \pm 0.09)M_{[8.0]_{\text{PAH}}}$ (fit only to $M_{[8.0]_{\text{PAH}}} < -16$); *Bottom panel:* $12 + \log \text{O/H} = (6.48 \pm 0.37) - (0.08 \pm 0.02)M_{[8.0]_{\text{PAH}}}$.

Fig. 9.— Integrated brightness of $[8.0]_{\text{PAH}}$ PAH emission, area-normalized star formation rate from Table 1, and oxygen abundance plotted as a function of the $[3.6] - [8.0]_{\text{PAH}}$ color. This color represents the amount of stellar emission relative to PAH emission. Dashed lines are fits to the data: *Top panel:* $M_{[8.0]_{\text{PAH}}} = (-16.55 \pm 0.65) - (2.52 \pm 0.52)([3.6] - [8.0]_{\text{PAH}})$; *Second panel:* $(U - B)_0 = (-0.33 \pm 0.05) + (-0.06 \pm 0.04)([3.6] - [8.0]_{\text{PAH}})$ (excluding HS 0822+3541); *Third panel:* $\log \text{SFR}/\text{area} = (-2.31 \pm 0.30) + (0.66 \pm 0.22)([3.6] - [8.0]_{\text{PAH}})$; *Bottom panel:* $12 + \log \text{O}/\text{H} = (7.71 \pm 0.18) + (0.27 \pm 0.13)([3.6] - [8.0]_{\text{PAH}})$.

Fig. 10.— Azimuthally-averaged $H\alpha$ emission relative to $[8.0]_{\text{PAH}}$ PAH emission for 4 of the galaxies with extended PAH emission. The galaxy profiles have been adjusted vertically to allow the different galaxies to be visually distinguished. NGC 6822 is not included because Milky Way cirrus prohibits quantitative information from the $8.0\ \mu\text{m}$ image.

Fig. 11.— False color images of the galaxies with extra-H II region $[8.0]_{\text{PAH}}$ PAH emission. $\text{H}\alpha$ is in green and the $[8.0]_{\text{PAH}}$ PAH image is in red. The field of view is $5.12'$ for IC 10, $2.08'$ for NGC 3738, $7.15'$ for NGC 4214, and $6.73'$ for NGC 4449. The NGC 6822 image is $14.40' \times 14.54'$. North is to the top and east to the left.

Fig. 12.— Deep displays of $H\alpha$ images of IC 10 and NGC 4449. The contours are of $[8.0]_{\text{PAH}}$ PAH emission. These images illustrate the PAH emission associated with gas shells and supergiant filaments. In NGC 4449 $H\alpha$ filaments 5 and 6 of Hunter & Gallagher (1992, 1997), also labelled here, have PAH emission associated with it in patches. There is also PAH emission associated with the supergiant H I shell of which filament 6 forms the ionized inner edge of a portion. In IC 10 H I PAH emission delineates shells number 5 and 7 of Wilcots & Miller (1998), labeled here near the centers of the holes.

Table 1. Galaxy Sample

Galaxy	Other Names ^a	D ^b (Mpc)	M_V^c	$(U - B)_0^d$	$(B - V)_0^d$	O/H ^e	$\log \dot{M}_D^f$ ($M_\odot$ /yr/kpc ²)	$\mu_0^V(\text{in})^g$ (mag arcsec ⁻²)	$R_D^V(\text{in})^g$ (arcmin)	$\mu_0^V(\text{out})^g$ (mag arcsec ⁻²)	$R_D^V(\text{out})^g$ (arcmin)
Im Galaxies											
CVnIdwA	UGCA 292	4.1	-12.65	-0.54	0.21	...	-2.64	24.30	0.54	...	...
DDO 50.....	PGC 23324, UGC 4305, HoII, VIIZw 223	3.4	-16.61	-0.41	0.22	7.83	-1.83	22.19	1.11	...	...
DDO 53.....	PGC 24050, UGC 4459, VIIZw 238	3.6	-13.84	-0.53	0.41	7.79	-2.50	23.81	0.69	...	...
DDO 63.....	PGC 27605, UGC 5139, HoI	3.8	-14.73	-0.18	0.20	...	-3.44	24.17	2.72	...	...
DDO 69.....	PGC 28868, UGC 5364, Leo A, Leo III	0.8	-11.67	-0.29	0.29	7.27	-3.00	...	...	23.01	0.81
DDO 75.....	PGC 29653, UGCA 205, Sextans A	1.3	-13.91	-0.54	0.19	7.49	-1.40	...	...	20.40	0.59
DDO 154.....	PGC 43869, UGC 8024, NGC 4789A	4.3	-14.51	-0.32	0.30	7.70	-2.60	23.06	0.55	...	...
DDO 155.....	PGC 44491, UGC 8091, GR 8	2.2	-12.53	-0.45	0.28	7.68	-1.50	21.72	0.23	...	...
DDO 165.....	PGC 45372, UGC 8201, IIZw 499	4.8	-15.69	-0.21	0.23	...	-3.52	23.32	1.69	21.72	0.50
DDO 187.....	PGC 50961, UGC 9128	2.5	-12.95	-0.18	0.30	7.62	-2.64	22.05	0.28	...	...
DDO 210.....	PGC 65367, Aquarius Dwarf	0.9	-10.88	-0.09	0.50	...	...	23.77	0.63	...	...
DDO 216.....	PGC 71538, UGC 12613, Pegasus Dwarf	0.9	-13.29	-0.03	0.66	7.93	-4.15	23.44	1.67	21.26	1.03
IC 10.....	PGC 1305, UGC 192	1.0	-17.11	...	0.59	8.18	-1.31	19.38	1.27	19.85	1.94
M81dwA	PGC 23521	3.6	-11.73	-0.33	0.27	...	...	23.32	0.25	...	...
NGC 3738....	PGC 35856, UGC 6565, Arp 234	4.9	-17.12	-0.16	0.42	8.36	-1.74	19.64	0.27	21.40	0.55
NGC 4214....	PGC 39225, UGC 7278	2.9	-17.56	-0.38	0.37	8.43	-1.10	20.42	0.86	...	...
NGC 4449....	PGC 40973, UGC 7592	3.9	-18.31	-0.44	0.36	8.32	-0.67	19.37	0.57	20.63	0.81
NGC 6822....	PGC 63616, IC 4895, DDO 209	0.5	-15.22	-0.34	0.46	8.27	-1.96	21.34	3.92	...	...
BCD Galaxies											
HS 0822+3542		10.9	-13.15	-0.90	0.26	7.35	-0.73	20.55	0.04	...	...
NGC 1705....	PGC 16282	5.0	-16.25	-0.52	0.28	8.21	-0.82	...	0.28	...	...
VIIZw 403 ...	PGC 35286, UGC 6456	4.4	-14.27	-0.49	0.28	7.70	-1.82	22.11	0.28	23.12	0.41

Table 1—Continued

Galaxy	Other Names ^a	D ^b (Mpc)	M_V ^c	$(U - B)_0$ ^d	$(B - V)_0$ ^d	O/H ^e	$\log \dot{M}_D$ ^f ($M_\odot$ /yr/kpc ²)	$\mu_0^V(\text{in})$ ^g (mag arcsec ⁻²)	$R_D^V(\text{in})$ ^g (arcmin)	$\mu_0^V(\text{out})$ ^g (mag arcsec ⁻²)	$R_D^V(\text{out})$ ^g (arcmin)
Sm Galaxies											
NGC 3109	PGC 29128,UGCA 194,DDO 236	1.3	...	...	...	8.06	-2.39	...	3.27	...	...

^aSelected alternate identifications obtained from NED.

^bDistances were taken from Hunter & Elmegreen (2006), and were determined from variable stars or the tip of the Red Giant Branch, if available in the literature. Other distances were determined from the radial velocity relative to the Galactic standard of rest V_{GSR} (RC3) and a Hubble constant of 65 km s⁻¹Mpc⁻¹. References for individual distances are given by Hunter and Elmegreen.

^c M_V are from Hunter & Elmegreen (2006).

^d UBV colors are from Hunter & Elmegreen (2006) except NGC 1705 whose colors are from de Vaucouleurs et al. (1991).

^eOxygen abundance is 12+log(O/H). References are given in Hunter & Hoffman (1999) except DDO 69 and DDO 187 (Pilyugin 2001), DDO 154 (Kennicutt & Skillman 2001), DDO 216 (Skillman et al. 1997), NGC 3109 (Richer & McCall 1995), NGC 4449 (Talent 1980), HS 0822+3542 (Kniazev et al. 2000), and NGC 1705 (Lee & Skillman 2004).

^fIntegrated star formation rate normalized to the area of the galaxy within R_D^V , from Hunter & Elmegreen (2004).

^gFor surface brightness profiles with two parts, the first central surface brightness and disk scale-length are the fit to the inner part of the profile, and the second set of values are for the fit to the outer part of the profile, from Hunter & Elmegreen (2006). The NGC 3109 surface photometry was measured from an H α off-band image, NGC 1705 from the Digitized Sky Survey. The radius at which the inner profile makes the transition to the outer profile R_{Br} are given in Tables 2 and 3.

Table 2. Integrated Properties for Galaxies Observed in Four IRAC Passbands^a

Galaxy	$M_{[3.6]}$	$[3.6]-[4.5]_*$	$V-[3.6]$	$M_{[8.0]_{\text{PAH}}}$	$[5.8]_{\text{PAH}}-[8.0]_{\text{PAH}}$	$[3.6]-[8.0]_{\text{PAH}}$	$M_{[4.5]_{\text{dust}}}$	$\mu_0^{[3.6]}(\text{in})^{\text{b}}$ (mag arcsec ⁻²)	$R_D^{[3.6]}(\text{in})^{\text{b}}$ (arcmin)	$\mu_0^{[3.6]}(\text{out})^{\text{b}}$ (mag arcsec ⁻²)	$R_D^{[3.6]}(\text{out})^{\text{b}}$ (arcmin)	R_{Br}^{b} (arcmin)
Im Galaxies												
CVnIdwA	-14.95	-0.53	2.29	<-13.84	...	...	<-11.29	22.24	0.46	...	...	...
	0.03	0.17	0.09	...	...	...	...	0.42	0.23	...	...	...
DDO 50	-18.77	0.09	2.21	-18.99	0.97	0.22	-16.17	19.57	0.93	...	...	...
	0.00	0.00	0.00	0.04	0.06	0.04	0.04	0.26	0.09	...	...	...
DDO 53	-16.10	0.03	2.38	-16.90	0.72	0.81	-14.14	21.25	0.63	...	...	...
	0.02	0.06	0.02	0.19	0.31	0.19	0.17	0.23	0.08	...	...	...
DDO 63	-16.38	-0.03	2.07	-15.09	...	-1.29	-12.20	22.10	2.75	...	...	...
	0.01	0.02	0.02	0.89	...	0.89	0.95	0.13	1.38	...	...	...
DDO 154	-15.82	-0.07	1.66	<-13.61	...	...	<-10.92	21.37	0.54	...	...	...
	0.01	0.03	0.02	...	...	...	...	0.12	0.05	...	...	...
DDO 165	-17.47	0.12	1.92	-17.61	...	0.14	<-11.44	21.53	1.79	20.51	0.69	1.06
	0.00	0.01	0.01	0.15	...	0.15	...	0.04	0.19	0.10	0.03	...
DDO 187	-14.35	0.27	1.68	<-13.18	...	...	<-10.71	20.23	0.26	...	...	...
	0.01	0.03	0.01	...	...	...	...	0.45	0.06	...	...	...
IC 10	-19.83	-0.17	2.72	-21.87	1.54	2.03	-16.54	16.93	1.45	17.27	2.04	1.58
	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.16	0.07	0.11	...
M81dwA	-13.36	-0.37	1.63	-13.18	...	-0.18	<-10.31	21.62	0.26	...	...	...
	0.04	0.22	0.07	1.32	...	1.32	...	0.17	0.02	...	...	...
NGC 3738	-19.62	-0.04	2.50	-20.33	1.42	0.71	-15.12	17.50	0.29	19.91	1.04	0.88
	0.00	0.00	0.00	0.01	0.03	0.01	0.05	0.08	0.01	0.16	0.09	...
NGC 4214	-20.01	-0.07	2.54	-22.10	1.58	2.08	-16.52	17.23	0.60	...	...	...
	0.00	0.00	0.00	0.00	0.01	0.00	0.04	0.09	0.02	...	...	...
NGC 4449	-21.16	-0.11	2.88	-23.74	1.76	2.57	-18.44	16.76	0.60	...	...	2.19
	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.08	0.02	...	...	...
NGC 6822	-17.99	-0.06	6.61	...	...	...	-12.59	18.96	4.46	...	...	...
	0.00	0.00	0.00	...	...	...	0.06	0.07	0.56	...	...	...
BCD Galaxies												
HS 0822+3542	-14.65	-0.08	1.50	-15.16	...	0.51	-13.09	19.86	0.05	...	...	...
	0.09	0.25	0.09	0.76	...	0.76	0.77	0.14	0.00	...	...	...

Table 2—Continued

Galaxy	$M_{[3.6]}$	$[3.6]-[4.5]_*$	$V-[3.6]$	$M_{[8.0]_{\text{PAH}}}$	$[5.8]_{\text{PAH}}-[8.0]_{\text{PAH}}$	$[3.6]-[8.0]_{\text{PAH}}$	$M_{[4.5]_{\text{dust}}}$	$\mu_0^{[3.6]}(\text{in})^{\text{b}}$ (mag arcsec $^{-2}$)	$R_D^{[3.6]}(\text{in})^{\text{b}}$ (arcmin)	$\mu_0^{[3.6]}(\text{out})^{\text{b}}$ (mag arcsec $^{-2}$)	$R_D^{[3.6]}(\text{out})^{\text{b}}$ (arcmin)	R_{Br}^{b} (arcmin)
NGC 1705	-18.55	0.01	...	-19.19	1.56	0.64	-15.43	17.84	0.25	...	...	...
	0.00	0.00	...	0.02	0.05	0.02	0.03	0.15	0.01	...	...	...
VIIZw 403	-16.27	-0.15	2.01	-16.97	1.09	0.69	-13.64	20.26	0.30	21.03	0.43	0.79
	0.01	0.04	0.02	0.12	0.24	0.12	0.20	0.05	0.01	0.05	0.01	...

^aSecond line are the uncertainties for the quantities in the first line. Values preceeded by “<” are upper limits. Quantities $[3.6]$ and $[4.5]_*$ are stellar properties; $[4.5]_{\text{dust}}$, $[5.8]_{\text{PAH}}$, and $[8.0]_{\text{PAH}}$ are measured from star-subtracted images. Separation of the dust and stellar components is discussed in §2.

^bFor surface brightness profiles with two parts, the first central surface brightness and disk scale length are the fit to the inner part of the profile, and the second set of values are for the fit to the outer part of the profile. R_{Br} is the radius at which this break occurs in the optical from Hunter & Elmegreen (2006).

Table 3. Integrated Properties for Galaxies Observed in only [4.5] and [8.0]^a

Galaxy	$M_{[4.5]*}$	$V-[4.5]*$	$M_{[8.0]_{\text{PAH}}}$	$[4.5]*-[8.0]_{\text{PAH}}$	$\mu_0^{[4.5]}(\text{in})^b$ (mag arcsec ⁻²)	$R_D^{[4.5]}(\text{in})^b$ (arcmin)	$\mu_0^{[4.5]}(\text{out})^b$ (mag arcsec ⁻²)	$R_D^{[4.5]}(\text{out})^b$ (arcmin)	R_{Br}^b (arcmin)
Im Galaxies									
DDO 69..	-13.14	1.76	<-9.25	...	...	...	21.39	0.99	1.16
	0.01	0.01	...	...	...	...	0.62	0.35	...
DDO 75..	-15.44	1.67	<-10.92	...	...	...	18.53	0.55	1.77
	0.01	0.02	...	...	...	...	0.24	0.03	...
DDO 155.	-14.17	1.83	-14.36	0.19	1 ...	0.24	...	...	...
	0.01	0.01	0.21	0.21	0.51	0.06	...	...	...
DDO 210.	-12.60	1.91	<-9.86	...	21.94	0.70	...	...	...
	0.06	0.06	...	...	0.15	0.08	...	...	...
DDO 216.	-15.38	2.34	-13.88	-1.51	21.11	1.80	...	...	5.42
	0.01	0.01	0.30	0.30	0.01	0.02	...	...	...
Sm Galaxies									
NGC 3109	-17.68	-0.52	-17.59	-0.09	20.83	3.36	...	...	...
	0.00	0.00	0.05	0.05	0.06	0.19	...	...	...

^aSecond line are the uncertainties for the quantities in the first line. Values preceeded by “<” are upper limits. Quantities labeled [4.5]* are stellar properties; quantities labeled [8.0]_{PAH} are measured from star-subtracted images. Separation of the dust and stellar components is discussed in §2. However, for the galaxies observed only in the [4.5] and [8.0] passbands, the hot dust could not be separated from the stellar component using the technique described there and was assumed to be negligible.

^bFor surface brightness profiles with two parts, the first central surface brightness and disk scale length are the fit to the inner part of the profile, and the second set of values are for the fit to the outer part of the profile. R_{Br} is the radius at which this break occurs in the optical from Hunter & Elmegreen (2006).

Table 4. Photometry of 3.6 μm excess Regions^a

Galaxy	Region ^b	R.A. (J2000.0)	Decl. (J2000.0)	R^c (pc)	M_V	$B - V$	$U - B$	$M_{[3.6]}$	$V - [3.6]$	$[3.6] - [4.5]_*^d$	$\mu_{3.6}^{Vega}$ ($L_{3.6}^{Vega} \text{ pc}^{-2}$)
IC 10	1	00 20 27.8	59 17 09	16.5	-5.3	1.0	...	-12.2	6.9	-0.07	150
					0.2	0.3	...	0.0	0.2	0.01	1
	2	00 20 27.9	59 17 05	16.5	-2.7	3.0	...	-11.8	9.1	-0.08	110
					1.9	9.9	...	0.0	1.9	0.01	1
	3	00 20 27.1	59 17 06	16.5	-5.2	0.4	...	-12.5	7.3	-0.07	200
					0.3	0.3	...	0.0	0.3	0.01	1
	4	00 20 28.8	59 16 56	16.5	...	...	...	-11.9	...	0.13	110
					...	...	...	0.0	...	0.01	1
	D1	00 20 28.0	59 17 01	43	...	...	...	-13.6	...	-0.10	80
					...	...	...	0.0	...	0.01	1
NGC 3738	1	11 35 48.6	54 31 32	107.8	-9.8	1.4	-2.9	-14.5	5.0	-0.04	30
					0.8	2.6	2.0	0.0	0.8	0.02	0
NGC 4214	1	12 15 40.9	36 19 04	47.3	-11.4	0.5	-1.0	-14.2	2.8	-0.07	120
					0.0	0.0	0.0	0.0	0.0	0.01	0
	2	12 15 40.8	36 19 10	35.4	-11.0	0.4	-1.0	-13.8	2.8	-0.07	140
					0.0	0.0	0.0	0.0	0.0	0.01	1
	D1	12 15 40.8	36 18 59	29.5	-7.7	0.4	-1.0	-12.2	4.5	-0.19	46
					0.1	0.1	0.1	0.0	0.1	0.03	0
	D2	12 15 36.8	36 20 02	84	-11.6	0.4	-0.1	-14.4	2.8	-0.03	46
					0.0	0.0	0.0	0.0	0.0	0.01	0
	1	12 28 10.7	44 05 34	82	...	...	...	-16.9	...	-0.10	470
					...	...	...	0.0	...	0.01	1
NGC 4449	D1	12 28 13.5	44 05 27	161	-14.0	0.4	-0.3	-16.8	2.7	-0.11	110
					0.0	0.0	0.0	0.0	0.0	0.01	0
	D2	12 28 08.7	44 05 20	136	-13.8	0.4	-0.3	-16.6	2.8	-0.11	120
					0.0	0.0	0.0	0.0	0.0	0.01	0
	D3	12 28 14.6	44 07 17	163	-12.7	0.2	-1.1	-15.8	3.1	-0.11	45
					0.0	0.0	0.0	0.0	0.0	0.02	0

^aSecond line are the uncertainties for the quantities in the first line.

^bRegions with “D” in their name are diffuse in the 3.6 μm image.

^cRadius of photometric aperture or, for polygons, the radius of the circle with the same area as the polygon.

^d[4.5]_{*} is the stellar contribution to the 4.5 μm image. Separation of the dust and stellar components is discussed in §2.

This figure "f1a.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f2.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f3a.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f3b.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f4.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f5.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f6.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f7.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f8.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>

This figure "f9.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/0604098v1>